

PRO- p LINK GROUPS AND p -HOMOLOGY GROUPS

JONATHAN HILLMAN¹, DANIEL MATEI² AND MASANORI MORISHITA³

1. School of Mathematics and Statistics, The University of Sydney, Sydney, NSW 2006, Australia
2. Graduate School of Mathematical Sciences, University of Tokyo, 3-8-1 Komaba, Meguro, Tokyo 153-8914, Japan
3. Department of Mathematics, Kanazawa University, Kakuma-machi, Kanazawa, Ishikawa, 920-1192, Japan

Contents

Introduction

1. Pro- p completion of a link group
2. p -adic Milnor invariants
3. Completed Alexander modules
4. Galois module structure of the p -homology group of a p -fold cyclic branched cover
5. Iwasawa type formulas for the p -homology groups of p^m -fold cyclic branched covers

Introduction

Let L be a tame link in the 3-sphere S^3 consisting of n knots $K_1, \dots, K_n$ and let G_L be the link group $\pi_1(X_L)$, $X_L = S^3 \setminus L$. For a prime number p , let $\widehat{G}_L$ denote the pro- p completion of the group G_L , $\widehat{G}_L = \varprojlim G_L/N$ where N runs over normal subgroups of G_L having p -power indices. By a theorem of J. Milnor [Mi], it is shown that $\widehat{G}_L$ has the following simple presentation as a pro- p group

$$\widehat{G}_L = \langle x_1, \dots, x_n \mid [x_1, y_1] = \dots = [x_n, y_n] = 1 \rangle$$

where x_i and y_i represent the meridian and longitude around K_i respectively (Theorem 1.2.1). The purpose of this paper is to use the pro- p link group $\widehat{G}_L$ and the associated group-theoretic invariants for the study of the p -homology groups of p^m -fold cyclic branched covers of S^3 along L , following the analogies between link theory and number theory [Mo1~4], [Rez1,2]. The invariants we derive from $\widehat{G}_L$ are the p -adic Milnor invariants and the completed Alexander module over the formal power series ring $\widehat{\Lambda}_n = \mathbf{Z}_p[[X_1, \dots, X_n]]$ with coefficients in the ring $\mathbf{Z}_p$ of p -adic integers. The tool involved here is the Fox differential calculus on a free pro- p group [Ih]. Although these invariants are simply p -adic analogues of the usual Milnor invariants and Alexander modules, it is natural to work over $\widehat{\Lambda}_n$ since the completed Alexander module can be presented over $\widehat{\Lambda}_n$ by a sort of

universal p -adic higher linking matrix $\widehat{T}_L$, called the *p -adic Traldi matrix*. This is defined in terms of the p -adic Milnor numbers and we can derive from $\widehat{T}_L$ systematically the “ p -primary” information on the homology of p^m -fold branched covers of L . This is an idea analogous to Iwasawa theory [Iw] which may also be regarded as a p -adic strengthening of the method employed by W. Massey [Mas] and L. Traldi [T]. We note that the method using the truncated Traldi matrices was considered in [Mat] to study the homology of unbranched covers.

The homology of cyclic branched covers of a link L is one of the basic invariants of L and has been extensively investigated by many authors. The Betti number and the order have been determined in terms of the Alexander (Hosokawa) polynomial ([HK],[MM],[S1] etc) and further the (Galois) module structure has been studied ([Da],[HS],[S2] etc), however most results are concerned mainly with the part which is prime to the covering degree. In [Rez1,2], A. Reznikov studied the p -homology of p -fold branched covers after the model of the classical problem on p -ideal class groups in number theory (see also [Mo1]). In this paper, we push this line of study in *arithmetic topology* further and determine the Galois module structure of the p -homology of a p -fold branched cover along a link completely in terms of the p -adic higher linking matrices. To be precise, let M be the p -fold cyclic branched cover of S^3 along L obtained from the completion of the p -fold total linking cover of X_L and let σ denote a generator of the Galois group of M over S^3 . The homology group $H_1(M, \mathbf{Z}_p) = H_1(M, \mathbf{Z}) \otimes_{\mathbf{Z}} \mathbf{Z}_p$ is then a module over the complete discrete valuation ring $\widehat{\mathcal{O}} := \mathbf{Z}_p[\langle \sigma \rangle] / (\sigma^{p-1} + \cdots + \sigma + 1) = \mathbf{Z}_p[\zeta]$, $\zeta := \sigma \bmod (\sigma^{p-1} + \cdots + \sigma + 1)$. Assume that $H_1(M, \mathbf{Z})$ is finite. Then the p -primary part $H_1(M, \mathbf{Z}_p)$ has $\mathfrak{p}$ -rank $n - 1$ ([Mo1],[Rez2]) so that it has form

$$H_1(M, \mathbf{Z}_p) = \bigoplus_{i=1}^{n-1} \widehat{\mathcal{O}} / \mathfrak{p}^{a_i} \quad (a_i \geq 1)$$

as $\widehat{\mathcal{O}}$ -module where $\mathfrak{p} := (\zeta - 1)$ is the maximal ideal of $\widehat{\mathcal{O}}$. Hence the determination of the Galois module structure of $H_1(N, \mathbf{Z}_p)$ is equivalent to that of the $\mathfrak{p}^k$ -rank

$$e_k := \#\{i \mid a_i \geq k\} \quad (k \geq 1).$$

Our main result is to give formulas for e_k ’s in terms of the higher linking matrices obtained by specializing the truncated p -adic Traldi matrices at $X_1 = \cdots = T_n = \zeta - 1$ (Theorem 4.1.3). For the simplest case of $k = 2$, our formula reads

$$e_2 = n - 1 - \text{rank}_{\mathbf{F}_p}(C \bmod p)$$

where $C = (C_{ij})$ is the linking matrix defined by $C_{ii} = -\sum_{j \neq i} \text{lk}(K_i, K_j)$ and $C_{ij} = \text{lk}(K_i, K_j)$ for $i \neq j$. In view of the analogy between the linking number and the power

residue symbols [Mo2,3], this is seen as a link-theoretic analog of L. Rédei's formula for the 4-rank of the class group of a quadratic field ([Réd1]), and our general result was partly suggested by the relation between Rédei's triple symbol and the 8-rank of a class group [Réd2]. In fact, the whole argument here can be translated into arithmetic [Mo5]. In the last section, we study the asymptotic behavior of the order $|H_1(M_m, \mathbf{Z}_p)|$ for the p^m -fold cyclic branched cover M_m as $m \rightarrow \infty$, following Iwasawa theory on $\mathbf{Z}_p$ -extensions [Iw]. Though our results obtained in this paper are rather elementary, they seem to indicate further possibilities of our arithmetic approach to link theory.

Acknowledgement. We would like to thank K. Murasugi for helpful communication.

Notation. Throughout this paper, we fix a prime number p . We denote by $\mathbf{F}_p$ the field with p elements and by $\mathbf{Z}_p$ the ring of p -adic integers. Let ord_p denote the additive p -adic valuation extended on the algebraic closure $\overline{\mathbf{Q}_p}$ of the p -adic field $\mathbf{Q}_p$ with $\text{ord}_p(p) = 1$ and set $|x|_p = p^{-\text{ord}_p(x)}$, $x \in \overline{\mathbf{Q}_p}$. We use the letter q to denote p or 0. For a topological (possibly discrete) group G , we denote by $G^{(k,q)}$ the k -th term of lower central q -series defined by $G^{(1,q)} = G$, $G^{(k+1,q)} = (G^{(k,q)})^q [G^{(k,q)}, G]$ where for closed subgroups A, B of G , $[A, B]$ stands for the closed subgroup of G generated by $[a, b] = aba^{-1}b^{-1}$, $a \in A, b \in B$. We simply write $G^{(k)}$ for $G^{(k,0)}$, the k -th term of the lower central series of G . For a pro-finite group G and a complete local ring R , we denote by $R[[G]]$ the completed group ring of G over R [Ko, §7].

1. Pro- p completion of a link group

1.1. *The pro- p completion of a link group.* Let L be a tame link in the 3-sphere S^3 consisting of n component knots $K_1, \dots, K_n$ and let G_L be the link group $\pi_1(X_L)$, $X_L = S^3 \setminus L$. After the work of K.T. Chen, J. Milnor [Mi] derived the following information about the presentation of the nilpotent quotient $G_L/G_L^{(k,q)}$. Let F be the free group on the n words $x_1, \dots, x_n$ where x_i represents the meridian m_i around K_i and let $\pi : F \rightarrow G_L$ be the meridional homomorphism defined by $\pi(x_i) = m_i$ ($1 \leq i \leq n$).

Theorem 1.1.1 ([Mi]). *For each $k \geq 1$ and i ($1 \leq i \leq n$), there is a word $y_i^{(k)}$ in $x_1, \dots, x_n$ representing the image of the i -th longitude in the quotient $G_L/G_L^{(k,q)}$ such that*

$$(1.1.2) \quad y_i^{(k)} \equiv y_i^{(k+1)} \mod F^{(k,q)}$$

and such that $\pi : F \rightarrow G_L$ induces the isomorphism

$$(1.1.3) \quad F/N_k F^{(k,q)} \simeq G_L/G_L^{(k,q)}$$

where N_k is the subgroup of F generated normally by $[x_1, y_1^{(k)}], \dots, [x_n, y_n^{(k)}]$.

Let $\widehat{G}_L$ be the pro- p completion of G_L , namely the inverse limit $\varprojlim G_L/N$ of the tower of quotients G_L/N which are finite p -groups. Since the quotients by the lower central p -series of G_L are cofinal in this tower, we have

$$\widehat{G}_L = \varprojlim_k G_L/G_L^{(k,p)}.$$

Since $\{y_i^{(k)} F^{(k,p)}\}_{k \geq 1}$ forms an inverse system in the tower $\{F/F^{(k,p)}\}_{k \geq 1}$ by (1.1.2), we define the pro- p word y_i to be $(y_i^{(k)} F^{(k,p)})$ in the free pro- p group $\widehat{F} := \varprojlim F/F^{(k,p)}$ which represents the i -th “longitude” in $\widehat{G}_L$ under the map $\widehat{\pi} : \widehat{F} \rightarrow \widehat{G}_L$ induced by π . By taking the inverse limit with respect to k in the isomorphism (1.1.3) of finite p -groups for $q = p$, we have the following

Theorem 1.1.2. *The map $\widehat{\pi}$ induces the isomorphism of pro- p groups*

$$\widehat{F}/\widehat{N} \simeq \widehat{G}_L$$

where $\widehat{N}$ is the closed subgroup of $\widehat{F}$ generated normally by $[x_1, y_1], \dots, [x_n, y_n]$. In particular, we have $\widehat{G}_K \simeq \mathbf{Z}_p$ for a knot K .

Remark 1.1.3. (1) By the construction above, we note $y_i \equiv y_i^{(k)} \pmod{\widehat{F}^{(k,q)}}$ (F is embedded in $\widehat{F}$).

(2) In view of the analogy between knots and primes, the pro- p link group $\widehat{G}_L$ is regarded as an analog of the maximal pro- p Galois group over the rational number field $\mathbf{Q}$ unramified outside prime numbers $p_1, \dots, p_n$, $p_i \equiv 1 \pmod p$ [Mo2].

Theorem 1.1.2 tells us that from the group-theoretic point of view, any link of n components looks like a pure braid link with n strings after the pro- p completion. In particular, by applying the method of D. Anick [A] to determine the graded quotients of the lower central series of a pure braid link group to our pro- p link group $\widehat{G}_L$, we see that the pro- p analog of Murasugi’s conjecture holds (cf. [L]). We define the *mod p linking diagram* of L to be the graph with vertices the components of L and an edge joining K_i and K_j if and only if the linking number $\text{lk}(K_i, K_j) \not\equiv 0 \pmod p$.

Theorem 1.1.4. *If the mod p linking diagram of L is connected, we have the isomorphisms*

$$\widehat{G}_L^{(q)} / \widehat{G}_L^{(q+1)} \simeq \widehat{F}_1^{(q)} / \widehat{F}_1^{(q+1)} \times \widehat{F}_{n-1}^{(q)} / \widehat{F}_{n-1}^{(q+1)} \quad \text{for } q \geq 1,$$

where $\widehat{F}_r$ denotes the free pro- p group of rank r .

1.2. *The p -goodness of a link group.* Let G be a group and $\widehat{G}$ be the pro- p completion of G . We then call G p -good if the natural map $G \rightarrow \widehat{G}$ induces the isomorphisms on cohomology $H^q(\widehat{G}, M) \xrightarrow{\sim} H^q(G, M)$ for all $q \geq 0$ for any finite p -primary $\widehat{G}$ -modules M (cf. [Se]).

Theorem 1.2.1. *A link group G_L is p -good.*

Proof. We shall say that a subgroup G of finite index in G_L is *open* if $[G_L : G]$ is a power of p . Let M be a finite p -primary $\widehat{G}_L$ -module. We shall show by induction on the length of M that if G is an open subgroup of G_L then there is a smaller open subgroup G_1 such that restriction from $H^2(G, M)$ to $H^2(G_1, M)$ is trivial.

Suppose first that $M = \mathbf{F}_p$, with trivial G_L -action, and let $H^*(G)$ denote $H^*(G, \mathbf{F}_p)$, for ease of reading. Since $[G_L : G]$ is finite and $G_L/G_L^{(2)} \cong \mathbf{Z}^n$ there is an epimorphism $\tau : G \rightarrow C = \mathbf{Z}/p\mathbf{Z}$. Then $K = \text{Ker}(\tau)$ is another open subgroup of G_L . The Hochschild-Serre spectral sequence for G as an extension of C by K has E_2 term $E_2^{p,q} = H^p(C, H^q(K))$, r^{th} differential d_r of bidegree $(r, 1-r)$ and converges to $H^*(G)$. Since $H^p(K) = 0$ for $p > 2$ there are only three nonzero rows, and since $H^*(G) = 0$ for $* > 2$ we see that $d_3^{p,2}$ is an isomorphism for all $p \geq 1$. The spectral sequence is an algebra over the ring $H^*(C) = E_2^{*,0}$. Since C has cohomological period 2, the cup product with a generator of $H^2(C) \cong \mathbf{F}_p$ induces isomorphisms $\gamma_2^{p,q} : E_2^{p,q} \cong E_2^{p+2,q}$ such that $d_2^{p+2,q} \gamma_2^{p,q} = \gamma_2^{p+2,q-1} d_2^{p,q}$ for all $p, q \geq 0$. Therefore we have the isomorphisms $\text{Ker}(d_2^{p,q}) \simeq \text{Ker}(d_2^{p+2,q})$, $\text{Im}(d_2^{p,q}) \simeq \text{Im}(d_2^{p+2,q})$ for any $p, q \geq 0$. In particular we have the isomorphisms $\gamma_3^{0,2} : E_3^{0,2} = \text{Ker}(d_2^{0,2}) \cong E_3^{2,2} = \text{Ker}(d_2^{2,2})$ and $\gamma_3^{3,0} : E_3^{3,0} = E_2^{3,0}/\text{Im}(d_2^{1,1}) \cong E_3^{5,0} = E_2^{5,0}/\text{Im}(d_2^{3,1})$ with $d_3^{2,2} \circ \gamma_3^{0,2} = \gamma_3^{2,1} \circ d_3^{0,2}$. It follows that $d_3^{0,2}$ is also an isomorphism, and so $E_\infty^{0,2} = 0$. But the edge homomorphism from $H^2(G)$ to $H^2(K)$ factors through $E_\infty^{0,2} \leq E_2^{0,2} = H^2(K)^C$, and so is 0.

In general, M has a finite composition series whose factors are copies of the simple module $\mathbf{F}_p$. Suppose that M_1 is a maximal proper submodule of M , with quotient $M/M_1 \cong \mathbf{F}_p$. Restriction from G to K induces a homomorphism from the exact sequences of cohomology corresponding to the coefficient sequence $0 \rightarrow M_1 \rightarrow M \rightarrow \mathbf{F}_p \rightarrow 0$. The result for $\mathbf{F}_p$ implies that the image of $H^2(G; M)$ lies in the image of $H^2(K, M_1)$. By the hypothesis of induction we may assume the result is true for M_1 , and so there is an open subgroup $K_1 < K$ such that restriction from $H^2(K, M_1)$ to $H^2(K_1, M_1)$ is trivial. Hence restriction from $H^2(G, M)$ to $H^2(K_1, M)$ is also trivial. This establishes the inductive step.

In particular, restriction from $H^2(G_L, M)$ to $H^2(J, M)$ is trivial, for some open subgroup J , and so the result follows, as in Exercise 1 of Chapter I. §2.6 of [Se]. (This exercise is stated in terms of profinite completions, but extends easily to the pro- p case). $\square$

Since the cohomological dimension $cd(G_L) \leq 2$, with equality if and only if L is nontrivial,

Theorem 1.2.4 gives the corresponding bound for the pro- p completion $\widehat{G}_L$.

Corollary 1.2.2. *The cohomological p -dimension $cd_p(\widehat{G}_L) \leq 2$.*

If the Milnor invariants of L are all 0 mod p (cf. Section 2), then $\widehat{G}_L$ is a free pro- p group and so $cd_p(\widehat{G}_L)$ may be strictly less than $cd(G_L)$. In particular, this is so if L is a nontrivial knot.

2. p -adic Milnor invariants

2.1. *The pro- p Fox differential calculus.* Let $\widehat{F}$ be the pro- p completion of the free group F on n generators $x_1, \dots, x_n$. Y. Ihara [Ih] extended the Fox differential calculus on the abstract free group F ([F]) to that on $\widehat{F}$. The basic result is stated as the following

Theorem 2.1.1 ([Ih]). *There exists a unique continuous $\mathbf{Z}_p$ -homomorphism*

$$\partial_i = \frac{\partial}{\partial x_i} : \mathbf{Z}_p[[\widehat{F}]] \longrightarrow \mathbf{Z}_p[[\widehat{F}]]$$

for each i ($1 \leq i \leq n$) such that any element $\alpha \in \mathbf{Z}_p[[\widehat{F}]]$ is expressed uniquely in the form

$$\alpha = \epsilon(\alpha)1 + \sum_{i=1}^n \partial_i(\alpha)(x_i - 1)$$

where ϵ is the augmentation map $\mathbf{Z}_p[[\widehat{F}]] \rightarrow \mathbf{Z}_p$.

The higher order derivatives are defined inductively by

$$\partial_{i_1} \cdots \partial_{i_r}(\alpha) = \partial_{i_1}(\partial_{i_2} \cdots \partial_{i_r}(\alpha)).$$

Here are some basic rules (cf. [Ih,2]).

(2.1.2) 1. If one restricts ∂_i to $\mathbf{Z}[F]$ under the natural embedding $\mathbf{Z}[F] \rightarrow \mathbf{Z}_p[[\widehat{F}]]$, we get the usual Fox derivative on $\mathbf{Z}[F]$ ([F]).

2. $\partial_i(x_j) = \delta_{ij}$ (Kronecker delta).

3. $\partial_i(\alpha\beta) = \partial_i(\alpha)\epsilon(\beta) + \alpha\partial_i(\beta)$ ($\alpha, \beta \in \mathbf{Z}_p[[\widehat{F}]]$).

4. $\partial_i(f^{-1}) = -f^{-1}\partial_i(f)$ ($f \in \widehat{F}$).

5. For $f \in \widehat{F}$ and $a \in \mathbf{Z}_p$, $\partial_i(f^a) = b\partial_i(f)$, where b is any element of $\mathbf{Z}_p[[\widehat{F}]]$ such that $b(f-1) = f^a - 1$.
6. Let $\widehat{F}'$ be another free pro- p group on $x'_1, \dots, x'_m$ and let $\varphi : \widehat{F} \rightarrow \widehat{F}'$ be a continuous surjective homomorphism. Then one has $\partial'_i(\varphi(\alpha)) = \sum_{j=1}^n \varphi(\partial_j(\alpha))\partial'_i(\varphi(x_j))$, where $\partial'_i = \frac{\partial}{\partial x'_i}$, $\alpha \in \mathbf{Z}_p[[\widehat{F}]]$.
7. For $f \in \widehat{F}$, $\epsilon(\partial_i^r(f)) = \binom{a}{r}$ where $a = \epsilon(\partial_i(f))$ and $\binom{a}{r} = \frac{a(a-1)\cdots(a-r+1)}{r!} \in \mathbf{Z}_p$.

Let $\mathbf{Z}_p\langle\langle X_1, \dots, X_n \rangle\rangle$ be the formal power series ring over $\mathbf{Z}_p$ in non-commuting variables $X_1, \dots, X_n$ which is compact in the topology taking the ideals $I(r)$ of power series with homogeneous components of degree $\geq r$ as the system of neighborhood of 0. The pro- p Magnus embedding M of $\widehat{F}$ into $\mathbf{Z}_p\langle\langle X_1, \dots, X_n \rangle\rangle^\times$ is defined by

$$M(x_i) = 1 + X_i, \quad M(x_i^{-1}) = 1 - X_i + X_i^2 + \dots$$

and it is extended to give the isomorphism $\mathbf{Z}_p[[\widehat{F}]] \simeq \mathbf{Z}_p\langle\langle X_1, \dots, X_n \rangle\rangle$ of compact $\mathbf{Z}_p$ -algebras. The resulting expansion of $\alpha \in \mathbf{Z}_p[[\widehat{F}]]$ is given by the Fox derivatives:

$$(2.1.3) \quad \begin{aligned} M(\alpha) &= \epsilon(\alpha) + \sum_{I=(i_1 \cdots i_r)} \epsilon_I(\alpha) X_{i_1} \cdots X_{i_r}, \\ \epsilon_I(\alpha) &= \epsilon(\partial_{i_1} \cdots \partial_{i_r}(\alpha)), \quad I = (i_1 \cdots i_r). \end{aligned}$$

Finally, we recall the following fact ([Ko], 7.14): Let $\widehat{J}_q$ be the two-sided ideal of $\mathbf{Z}_p[[\widehat{F}]]$ generated by q and the augmentation ideal $I_{\mathbf{Z}_p[[\widehat{F}]]}$. Then for $f \in \widehat{F}$ and $k \geq 1$, we have

$$(2.1.4) \quad \begin{aligned} f \in \widehat{F}^{(k,q)} &\iff f - 1 \in \widehat{J}_q \\ &\iff M(f) = 1 + (\text{term of degree } \geq k) \quad (\text{for case } q = 0). \end{aligned}$$

2.2. p -adic Milnor invariants. We keep the same notation as in Section 1. Let $\widehat{F}$ be the free pro- p group on $x_1, \dots, x_n$ where each x_i represents the i -th meridian. For a multi-index $I = (i_1 \cdots i_r)$, $r \geq 1$, we define the p -adic Milnor number $\hat{\mu}(I)$ by

$$(2.2.1) \quad \begin{aligned} \hat{\mu}(I) &:= \epsilon_{I'}(y_{i_r}) \quad I' = (i_1 \cdots i_{r-1}) \\ &= \epsilon(\partial_{i_1} \cdots \partial_{i_{r-1}}(y_{i_r})) \end{aligned}$$

where $y_j \in \widehat{F}$ represents the j -th “longitude” in $\widehat{G}_L$ (cf. Section 1.1). By convention, we set $\hat{\mu}(I) = 0$ for $|I| = r = 1$. We let $\hat{\Delta}(I)$ denote the ideal of $\mathbf{Z}_p$ generated by $\hat{\mu}(J)$ where

J runs over all cyclic permutations of proper subsequences of I . We then define the p -adic Milnor invariant $\bar{\mu}(I)$ by

$$(2.2.2) \quad \bar{\mu}(I) := \hat{\mu}(I) \bmod \hat{\Delta}(I).$$

Since the usual Milnor number $\mu(I)$, $I = (i_1 \cdots i_r)$ is defined by $\epsilon(\partial_{i_1} \cdots \partial_{i_{r-1}}(y_{i_r}^{(r)}))$ by (2.1.1), 1, Remark 1.1.3, (1) and (2.2.1) yield

$$(2.2.3) \quad \hat{\mu}(I) = \mu(I) \text{ and } \hat{\Delta}(I) = \Delta(I)$$

as elements in $\mathbf{Z}_p$. Hence, we have

(2.2.4) $\bar{\mu}(I)$'s are isotopy invariants of L and satisfy the same properties such as the cycle symmetry and Shuffle relations $\bar{\mu}(I)$'s enjoy ([Mi]).

Theorem 1.1.2 and (2.1.4) implies the following:

(2.2.5) All $\bar{\mu}(I) = 0$ for $|I| < r$ if and only if $\hat{\pi} : \hat{F} \rightarrow \widehat{G}_L$ induces an isomorphism $\hat{F}/\hat{F}^{(r,q)} \simeq \widehat{G}_L/\widehat{G}_L^{(r,q)}$. In particular, if all p -adic Milnor invariants of L are zero, one has $\widehat{G}_L \simeq \hat{F}$. This is the case for boundary links.

Finally, we remark that (2.1.2), 7 implies

$$(2.2.6) \quad \hat{\mu}(\underbrace{i \cdots i}_r j) = \binom{\text{lk}(K_i, K_j)}{r} \text{ for } i \neq j.$$

The Milnor invariant is also given by the Massey products in the cohomology of $\widehat{G}_L$. For the normalized Massey system in profinite group cohomology and sign convention, we refer to [Mo3]. Let $\xi_1, \dots, \xi_n$ be the $\mathbf{Z}_p$ -basis of $H^1(\widehat{G}_L, \mathbf{Z}_p)$ dual to the meridians m_i 's, and let $\eta_j \in H_2(\widehat{G}_L, \mathbf{Z}_p)$ be the image of $[x_j, y_j]$ under the transgression $H_1(\hat{N}, \mathbf{Z}_p)_{\widehat{G}_L} \rightarrow H_2(\widehat{G}_L, \mathbf{Z}_p)$. Then for $I = (i_1 \cdots i_r)$, $r \geq 2$, there is a normalized Massey system M for the product $\langle \xi_{i_1}, \dots, \xi_{i_r} \rangle \in H_2(\widehat{G}_L, \mathbf{Z}_p/\hat{\Delta}(I))$ so that

$$(2.2.7) \quad \langle \xi_{i_1}, \dots, \xi_{i_r} \rangle(\eta_j) = \begin{cases} (-1)^r \bar{\mu}(I) & j = i_r \neq i_1, \\ (-1)^{r+1} \bar{\mu}(I) & j = i_1 \neq i_r, \\ 0 & \text{otherwise.} \end{cases}$$

3. Completed Alexander modules

3.1. *The Alexander module of $\widehat{G}_L$.* The Alexander module of a finitely presented pro- p group was introduced in several modes in [Mo2]. As a particular case, the Alexander module of the pro- p link group $\widehat{G}_L$ is defined using the pro- p Fox free differential calculus as follows. We keep the same notation as in Section 1 and 2. Let $\widehat{\psi}$ be the abelianization map $\widehat{G}_L \rightarrow \widehat{H} := \widehat{G}_L / \widehat{G}_L^{(2)} = \mathbf{Z}_p^n$ and denote by the same $\widehat{\psi}$ the continuous $\mathbf{Z}_p$ -homomorphism $\mathbf{Z}_p[[\widehat{G}_L]] \rightarrow \mathbf{Z}_p[[\widehat{H}]]$ on the completed group rings. We identify $\mathbf{Z}_p[[\widehat{H}]]$ with the commutative formal power series ring $\widehat{\Lambda}_n := \mathbf{Z}_p[[X_1, \dots, X_n]]$ over $\mathbf{Z}_p$ by setting $t_i := \widehat{\psi} \circ \widehat{\pi}(x_i) = 1 + X_i$. By Theorem 1.2.1, we call the matrix over $\widehat{\Lambda}_n$

$$(3.1.1) \quad \widehat{P}_L = \left(\widehat{\psi} \circ \widehat{\pi}(\partial_j([x_i, y_i])) \right)$$

the *Alexander matrix* of $\widehat{G}_L$. We then define the *Alexander module* $\widehat{A}_L$ of $\widehat{G}_L$ by the compact $\widehat{\Lambda}_n$ -module presented by $\widehat{P}_L$:

$$(3.1.2) \quad \widehat{A}_L := \text{Coker}(\widehat{\Lambda}_n^n \xrightarrow{\widehat{P}_L} \widehat{\Lambda}_n^n)$$

and call $\widehat{A}_L$ the *completed Alexander module* of L over $\widehat{\Lambda}_n$. Let $\widehat{\Lambda} = \mathbf{Z}_p[[X]]$, the *Iwasawa algebra* and $\tau : \widehat{\Lambda}_n \rightarrow \widehat{\Lambda}$ the reducing homomorphism defined by $\tau(X_i) = X$ ($1 \leq i \leq n$). Then the *reduced completed Alexander module* $\widehat{A}_L^{\text{red}}$ is defined by the compact $\widehat{\Lambda}$ -module

$$(3.1.3) \quad \widehat{A}_L^{\text{red}} := \tau(\widehat{A}_L) = \widehat{A}_L \otimes_{\widehat{\Lambda}_n} \widehat{\Lambda}$$

which is presented by $\tau(\widehat{P}_L) = \widehat{P}_L(X, \dots, X)$. For a knot K , $\widehat{P}_L = O_n$ (zero matrix) and so we have

$$(3.1.4) \quad \widehat{A}_L = \widehat{A}_L^{\text{red}} = \widehat{\Lambda}.$$

We also define the *i -th completed Alexander ideal* $\widehat{E}_i(L)$ of L by the i -th elementary ideal $E_i(\widehat{A}_L)$ of $\widehat{A}_L$ and *i -th p -adic Alexander series* $\widehat{\Delta}_i(L)$ of L by the greatest common divisor $\Delta_i(\widehat{E}_i(L))$ of generators of the ideal $\widehat{E}_i(L)$:

$$(3.1.5) \quad \widehat{E}_i(L) := E_i(\widehat{A}_L), \quad \widehat{\Delta}_i(L) := \Delta_i(\widehat{E}_i(L))$$

The relation with the usual Alexander module is given as follows. Let $\psi : G_L \rightarrow H = H_1(X_L, \mathbf{Z})$ be the abelianization map which induces the ring homomorphism $\psi : \mathbf{Z}[G_L] \rightarrow$

$\mathbf{Z}[H]$ on the group rings where $\mathbf{Z}[H]$ is identified with the Laurent polynomial ring $\Lambda_n = \mathbf{Z}[t_1^{\pm 1}, \dots, t_n^{\pm 1}]$, $t_i = \psi \circ \pi(x_i)$. Given a presentation $G_L = \langle x_1, \dots, x_m \mid r_1 = \dots r_l = 1 \rangle$ ($m \geq n$), the Alexander module of L is given as the Λ_n -module presented by the Alexander matrix $(\psi \circ \pi(\partial_j(r_i)))$. By Theorem 1.1.1, we can take $m = n$ and the relators to be $[x_i, y_i^{(k+1)}]$ ($1 \leq i \leq n$) and some finite number of generators $f_i^{(k+1)}$ of $F^{(k+1,p)}$ ($k \geq 1$). We let $J_p = \widehat{J}_p \cap \mathbf{Z}[F]$ and $J_{\Lambda_n, p} = \psi \circ \pi(J_p)$. Since $\partial_j(f_i^{(k+1)}) \in J_p^k$ by (2.1.4), passing to the quotients modulo $(J_{\Lambda_n, p})^k$, $A_L / (J_{\Lambda_n, p})^k A_L = A_L \otimes_{\Lambda_n} \Lambda_n / (J_{\Lambda_n, p})^k$ is presented by the matrix $(\psi \circ \pi(\partial_j([x_i, y_i^{(k+1)}])) \bmod (J_{\Lambda_n, p})^k)$. Here we see by Remark 1.1.3, (1) that the elements $\{\psi \circ \pi(\partial_j([x_i, y_i^{(k+1)}])) \bmod (J_{\Lambda_n, p})^k\}$ form an inverse system with respect to k and its limit is given as $\widehat{\psi} \circ \widehat{\pi}(\partial_j([x_i, y_i]))$ under the identification $\varprojlim_k \Lambda_n / (J_{\Lambda_n, p})^k = \widehat{\Lambda}_n$ defined by $t_i = 1 + X_i$. Hence by (3.1.2) we have

$$(3.1.6) \quad \widehat{A}_L = \varprojlim_k A_L \otimes_{\Lambda_n} (\Lambda_n / (J_{\Lambda_n, p})^k) = A_L \otimes_{\Lambda_n} \widehat{\Lambda}_n.$$

Similarly, $\widehat{A}_L^{red}$ is related with the usual reduced Alexander module $A_L^{red} = \tau(A_L)$ by

$$(3.1.7) \quad \widehat{A}_L^{red} = A_L \otimes_{\Lambda} \widehat{\Lambda}$$

where $\Lambda = \mathbf{Z}[t^{\pm 1}]$ is embedded into $\widehat{\Lambda}$ by $t = 1 + X$.

3.2. The p -adic Traldi matrix. The Alexander matrix $\widehat{P}_L$ of $\widehat{G}_L$ (3.1.1) is computed explicitly as a *universal p -adic higher linking matrix* in terms of p -adic Milnor numbers. This is regarded as a p -adic strengthening of the work by L. Traldi [Tr].

Definition 3.2.1. The p -adic Traldi matrix $\widehat{T}_L = (\widehat{T}_L(i, j))$ of L over $\widehat{\Lambda}_n$ is defined by

$$\widehat{T}_L(i, j) = \begin{cases} -\sum_{r \geq 1} \sum_{\substack{1 \leq i_1, \dots, i_r \leq n \\ i_r \neq i}} \hat{\mu}(i_1 \cdots i_r i) X_{i_1} \cdots X_{i_r} & i = j \\ \hat{\mu}(ji) X_i + \sum_{r \geq 1} \sum_{1 \leq i_1 \cdots i_r \leq n} \hat{\mu}(i_1 \cdots i_r ji) X_i X_{i_1} \cdots X_{i_r} & i \neq j. \end{cases}$$

and we also define the *reduced p -adic Traldi matrix* $\widehat{T}_L^{red}$ of L over $\widehat{\Lambda}$ by

$$\widehat{T}_L^{red} := \tau(\widehat{T}_L) = \widehat{T}_L(X, \dots, X).$$

Our theorem is then stated as

Theorem 3.2.2. *The p -adic Traldi matrix $\widehat{T}_L$ gives a presentation matrix for the completed Alexander module $\widehat{A}_L$ over $\widehat{\Lambda}_n$, and the reduced p -adic Traldi matrix $\widehat{T}_L^{red}$ gives a presentation matrix for the reduced completed Alexander module $\widehat{A}_L^{red}$ over $\widehat{\Lambda}$.*

Proof. By (3.1.1), (3.1.2) and (3.1.3), it suffices to show $\widehat{\psi} \circ \widehat{\pi}(\partial_j([x_i, y_i])) = \widehat{T}_L(i, j)$. By the rules 2 $\sim$ 4 of (2.1.2), we have

$$\partial_j([x_i, y_i]) = (1 - x_i y_i x_i^{-1}) \delta_{ij} + x_i (1 - y_i x_i^{-1} y_i^{-1}) \partial_j(y_i).$$

Here $y_i = 1 + \sum_{r \geq 1} \sum_{1 \leq i_1, \dots, i_r \leq n} \hat{\mu}(i_1 \cdots i_r i) (x_{i_1} - 1) \cdots (x_{i_r} - 1)$ by (2.1.3) and (2.2.1). Hence we get

$$(3) \quad \begin{aligned} \widehat{\psi} \circ \widehat{\pi}(\partial_j([x_i, y_i])) &= \delta_{i,j} - \sum_{r \geq 1} \sum_{1 \leq i_1, \dots, i_r \leq n} \hat{\mu}(i_1 \cdots i_r i) X_{i_1} \cdots X_{i_r} \\ &\quad + \hat{\mu}(ji) X_i + \sum_{r \geq 1} \sum_{1 \leq i_1, \dots, i_r \leq n} \hat{\mu}(i_1 \cdots i_r ji) X_i X_{i_1} \cdots X_{i_r}. \end{aligned}$$

which yields the assertion. $\square$

The following is an extension of (3.1.4).

Corollary 3.2.3. *For a link whose p -adic Milnor invariants are all zero, we have*

$$\widehat{A}_L \simeq \widehat{\Lambda}_n^n, \quad \widehat{A}_L^{red} \simeq \widehat{\Lambda}^n.$$

This is the case for boundary links.

Finally, we introduce the *truncated p -adic Traldi matrices*.

Definition 3.2.4. For $k \geq 2$, the k -th truncated p -adic Traldi matrix $\widehat{T}_L^{(k)} = (\widehat{T}_L^{(k)}(i, j))$ is defined by

$$\widehat{T}_L^{(k)}(i, j) = \begin{cases} -\sum_{r=1}^{k-1} \sum_{\substack{1 \leq i_1, \dots, i_r \leq n \\ i_r \neq i}} \hat{\mu}(i_1 \cdots i_r i) X_{i_1} \cdots X_{i_r} & i = j \\ \hat{\mu}(ji) X_i + \sum_{r=1}^{k-2} \sum_{1 \leq i_1, \dots, i_r \leq n} \hat{\mu}(i_1 \cdots i_r ji) X_i X_{i_1} \cdots X_{i_r} & i \neq j \end{cases}$$

and we also define the k -th truncated reduced p -adic Traldi matrix $\widehat{T}_L^{\text{red},(k)}$ by

$$\widehat{T}_L^{\text{red},(k)} := \tau(\widehat{T}_L^{(k)}) = \widehat{T}_L^{(k)}(X, \dots, X).$$

We note that $\widehat{T}_L^{\text{red},(2)}$ is the *linking matrix* multiplied by X , where the linking matrix $C = (C_{ij})$ is defined by $C_{ii} = -\sum_{j \neq i} \text{lk}(K_i, K_j)$ and $C_{ij} = \text{lk}(K_i, K_j)$ for $i \neq j$. Thus the p -adic Traldi matrix $\widehat{T}_L$ is regarded as a *universal higher linking matrix* over $\widehat{\Lambda}_n$ which contains all information on the completed Alexander module. In the following section, we derive from $\widehat{T}_L$ the information on the p -homology groups of p^m -fold cyclic branched covers along L .

4. Galois module structure for the p -homology group of a p -fold cyclic branched cover

4.1. *Galois module structure of the p -homology of a p -fold cover.* Let X_∞ be the infinite cyclic cover of X_L associated to the kernel of the homomorphism $G_L \rightarrow \langle t \rangle$ sending each meridian to t . Let M be the completion of the p -fold subcover of X_∞ over X_L so that M is a p -fold cyclic cover of S^3 branched along L . We set $\nu_d(t) = t^{d-1} + \dots + t + 1$ for $d \geq 1$. Let $\phi : M \rightarrow S^3$ be the covering map and σ a generator of its Galois group. Since $\nu_p(\sigma) = \text{tr} \circ \phi_* = 0$ where $\text{tr} : H_1(S^3, \mathbf{Z}) \rightarrow H_1(M, \mathbf{Z})$ is the transfer, we can regard $H_1(M, \mathbf{Z})$ as a module over the Dedekind ring $\mathcal{O} = \mathbf{Z}[\langle \sigma \rangle] / (\nu_p(\sigma)) = \mathbf{Z}[\zeta]$, $\zeta = \sigma \bmod (\nu_p(\sigma))$ is a primitive p -th root of 1. Hence $H_1(M, \mathbf{Z}_p) = H_1(M, \mathbf{Z}) \otimes_{\mathbf{Z}} \mathbf{Z}_p$ is regarded as a module over the complete discrete valuation ring $\widehat{\mathcal{O}} = \mathcal{O} \otimes_{\mathbf{Z}} \mathbf{Z}_p = \mathbf{Z}_p[\zeta]$. Note that $\widehat{\mathcal{O}}$ is the completion of $\mathcal{O}$ with respect to the maximal ideal $\mathfrak{p}$ generated by the prime element $\pi := \zeta - 1$ and the residue field $\widehat{\mathcal{O}}/\mathfrak{p}$ is $\mathbf{F}_p$. By Theorem 2.2.1, we can derive the following information on a presentation matrix for the $\widehat{\mathcal{O}}$ -module $H_1(M, \mathbf{Z}_p)$. Note that the evaluation of a power series $F(X) \in \mathbf{Z}_p[[X]]$ at $s = \pi$ makes sense in the $\mathfrak{p} = (\pi)$ -adically complete ring $\widehat{\mathcal{O}}$.

Theorem 4.1.1. *A presentation matrix for $H_1(M, \mathbf{Z}_p) \oplus \widehat{\mathcal{O}}$ over $\widehat{\mathcal{O}}$ is given by $\widehat{T}_L^{\text{red}}(\pi)$. Further, for any integer $k \geq 2$, a presentation matrix for $(H_1(M, \mathbf{Z}_p) \otimes_{\widehat{\mathcal{O}}} \widehat{\mathcal{O}}/\mathfrak{p}^k) \oplus \widehat{\mathcal{O}}/\mathfrak{p}^k$ over $\widehat{\mathcal{O}}/\mathfrak{p}^k$ is given by $\widehat{T}_L^{\text{red},(k)}(\pi)$. Here $\widehat{T}_L^{\text{red}}$ (resp. $\widehat{T}_L^{\text{red},(k)}$) is the reduced (resp. reduced truncated) Traldi matrix defined in Section 3.2.*

Proof. Note that the well-known relation ([S1, Theorem 6])

$$H_1(M, \mathbf{Z}) \simeq H_1(X_\infty, \mathbf{Z}) / \nu_p(t) H_1(X_\infty, \mathbf{Z})$$

is an $\widehat{\mathcal{O}}$ -isomorphism since σ acts on the r.h.s by t . Hence we have the following isomorphisms over $\widehat{\mathcal{O}}$

$$\begin{aligned} H_1(M, \mathbf{Z}_p) &\simeq (H_1(X_\infty, \mathbf{Z})/\nu_p(t)H_1(X_\infty, \mathbf{Z})) \otimes_{\mathbf{Z}} \mathbf{Z}_p \\ &\simeq H_1(X_\infty, \mathbf{Z}) \otimes_{\Lambda} (\Lambda \otimes_{\mathbf{Z}} \mathbf{Z}_p)/(\nu_p(t)) \\ &\simeq H_1(X_\infty, \mathbf{Z}) \otimes_{\Lambda} (\widehat{\Lambda}/(\nu_p(1+X))) \\ &\simeq H_1(X_\infty, \mathbf{Z}) \otimes_{\Lambda} \widehat{\mathcal{O}}. \end{aligned}$$

Since $A_L^{red} \simeq H_1(X_\infty, \mathbf{Z}) \oplus \Lambda$ as Λ -module ([H, 5.4]), tensoring with $\widehat{\mathcal{O}}$ over Λ , we have an isomorphism of $\widehat{\mathcal{O}}$ -modules

$$A_L^{red} \otimes_{\Lambda} \widehat{\mathcal{O}} \simeq H_1(M, \mathbf{Z}_p) \oplus \widehat{\mathcal{O}}.$$

Since the l.h.s is same as $\widehat{A}_L^{red} \otimes_{\widehat{\Lambda}} \widehat{\mathcal{O}}$, the first assertion follows from Theorem 3.2.4. The second assertion is obtained from the first one by taking modulo $\mathfrak{p}^k$. $\square$

Now, we assume that $H_1(M, \mathbf{Z})$ is finite so that $H_1(M, \mathbf{Z}_p)$ is the p -primary part of $H_1(M, \mathbf{Z})$. Using Theorem 3.1.1, we will see the $\widehat{\mathcal{O}}$ -module structure of $H_1(M, \mathbf{Z}_p)$ more precisely. First, we recall the following result on the $\mathfrak{p}$ -rank of $H_1(M, \mathbf{Z}_p)$ (cf. [Mo1],[Rez]).

Lemma 4.1.2. $H_1(M, \mathbf{Z}_p) \otimes_{\widehat{\mathcal{O}}} \mathbf{F}_p$ has dimension $n - 1$ over $\mathbf{F}_p$.

Proof. By [Mo1], the map $\Phi : H_1(M, \mathbf{Z}) \rightarrow \mathbf{F}_p^n$ defined by $\Phi(c) := (\text{lk}(\phi_*(c), K_i) \bmod p)$ induces an isomorphism

$$H_1(M, \mathbf{Z})/(\sigma - 1)H_1(M, \mathbf{Z}) \simeq \{(\xi_i) \in \mathbf{F}_p^n \mid \sum_{i=1}^n \xi_i = 0\} \simeq \mathbf{F}_p^{n-1}$$

where the l.h.s is $H_1(M, \mathbf{Z}) \otimes_{\mathcal{O}} \mathcal{O}/\mathfrak{p} = H_1(M, \mathbf{Z}_p) \otimes_{\widehat{\mathcal{O}}} \mathbf{F}_p$, and hence we are done. $\square$

By Lemma 4.1.2, $H_1(M, \mathbf{Z}_p)$ has the form

$$H_1(M, \mathbf{Z}_p) = \bigoplus_{i=1}^{n-1} \widehat{\mathcal{O}}/\mathfrak{p}^{a_i} \quad (a_i \geq 1)$$

as $\widehat{\mathcal{O}}$ -module. Hence, the determination of $\widehat{\mathcal{O}}$ -module structure of $H_1(M, \mathbf{Z}_p)$ is equivalent to the determination of the $\mathfrak{p}^k$ -rank

$$e_k := \#\{i \mid a_i \geq k\} = \dim_{\mathbf{F}_p} H_1(M, \mathbf{F}_p) \otimes_{\widehat{\mathcal{O}}} (\mathfrak{p}^{k-1}/\mathfrak{p}^k) \quad (k \geq 1).$$

We describe the $\mathfrak{p}^k$ -rank e_k in terms of $\widehat{T}_L^{red,(k)}(\pi)$. For an n by n matrix A over $\widehat{\mathcal{O}}$, we denote by $A \otimes_{\widehat{\mathcal{O}}} (\mathfrak{p}^{k-1}/\mathfrak{p}^k)$ the $\mathbf{F}_p$ -linear map on the $\mathbf{F}_p$ -vector space $(\mathfrak{p}^{k-1}/\mathfrak{p}^k)^n$ induced by A . Then Theorem 4.1.1 is restated as

Theorem 4.1.3. *Notation and assumption being as above, we have*

$$e_k = n - 1 - \text{rank}_{\mathbf{F}_p}(\widehat{T}_L^{red,(k)}(\pi) \otimes_{\widehat{\mathcal{O}}} (\mathfrak{p}^{k-1}/\mathfrak{p}^k)), \quad k \geq 2.$$

Here we may call $\widehat{T}_L^{red,(k)}(\pi)$ the k -th higher linking matrix in view of the following

Corollary 4.1.4. *For $k = 2$, we have*

$$e_2 = n - 1 - \text{rank}_{\mathbf{F}_p}(C \bmod p)$$

where $C = (C_{ij})$ is the linking matrix of L defined by $C_{ii} = -\sum_{j \neq i} \text{lk}(K_i, K_j)$ and $C_{ij} = \text{lk}(K_i, K_j)$ for $i \neq j$.

Proof. In fact, we have, by definition,

$$\widehat{T}_L^{red,(2)}(\pi) = \pi C$$

and hence $\text{rank}_{\mathbf{F}_p}(\widehat{T}_L^{red,(2)} \otimes_{\widehat{\mathcal{O}}} \mathfrak{p}/\mathfrak{p}^2) = \text{rank}_{\mathbf{F}_p}(C \bmod p)$. $\square$

4.2. 2-component case. We suppose $n = 2$ and keep to assume $H_1(M, \mathbf{Z})$ is finite. By Lemma 4.1.2, $H_1(M, \mathbf{Z}_p)$ has the $\mathfrak{p}$ -rank 1 so that we have

$$H_1(M, \mathbf{Z}_p) = \widehat{\mathcal{O}}/\mathfrak{p}^a, \quad a \geq 1.$$

Hence $e_k = 0$ or 1 for $k \geq 2$, and by Theorem 4.1.3 we have

$$e_k = 1 \iff \widehat{T}_L^{red,(k)}(\pi) \equiv O_2 \bmod \pi^k.$$

Since $\widehat{T}_L^{red,(k)}(1, 2)(\pi) = -\widehat{T}_L^{red,(k)}(1, 1)(\pi)$, $\widehat{T}_L^{red,(k)}(2, 2)(\pi) = -\widehat{T}_L^{red,(k)}(2, 1)(\pi)$, we have the following

Theorem 4.2.1. *Suppose $n = 2$. For each $k \geq 1$, assuming $e_k = 1$, we have*

$$e_{k+1} = 1 \iff \begin{cases} \sum_{r=1}^k \sum_{i_1, \dots, i_{r-1}=1,2} \hat{\mu}(i_1 \cdots i_{r-1} 21) \pi^r \equiv 0 \bmod \pi^{k+1}, \\ \sum_{r=1}^k \sum_{i_1, \dots, i_{r-1}=1,2} \hat{\mu}(i_1 \cdots i_{r-1} 12) \pi^r \equiv 0 \bmod \pi^{k+1}. \end{cases}$$

We give the condition in Theorem 4.2.1 in more concise forms for lower k . In the following computation, we use simply the usual Milnor number $\mu(I)$ instead of $\hat{\mu}(I)$ by (2.2.3).

Example 4.2.2. e_2 : Since $e_1 = 1$, we have by Theorem 4.2.1

$$(4.2.2.1) \quad e_2 = 1 \iff \mu(12)\pi \equiv 0 \pmod{\pi^2} \iff \text{lk}(K_1, K_2) \equiv 0 \pmod{p}.$$

e_3 : Assume $\text{lk}(K_1, K_2) \equiv 0 \pmod{p}$. By Theorem 4.2.1, we have

$$e_3 = 1 \iff \begin{cases} \mu(21)\pi + (\mu(121) + \mu(221))\pi^2 \equiv 0 \pmod{\pi^3}, \\ \mu(12)\pi + (\mu(112) + \mu(212))\pi^2 \equiv 0 \pmod{\pi^3}. \end{cases}$$

By cycle symmetry, $\mu(121) \equiv \mu(112)$, $\mu(221) \equiv \mu(212) \pmod{\mu(12)}$. Here $\mu(112) \equiv \mu(221) \equiv \binom{\mu(12)}{2} \pmod{\mu(12)}$ by (2.2.6). Thus we have $\mu(121) + \mu(221) \equiv \mu(112) + \mu(212) \equiv \mu(12) \equiv 0 \pmod{p}$. Hence, we have

$$(4.2.2.2) \quad e_3 = 1 \iff \text{lk}(K_1, K_2) \equiv 0 \pmod{p^2}.$$

As one easily see, this condition is also equivalent to

$$(4.2.2.3) \quad \mu(112) \equiv \mu(221) \equiv 0 \pmod{p}.$$

e_4 : Assume $\text{lk}(K_1, K_2) \equiv 0 \pmod{p^2}$. By Theorem 4.2.1, we have

$$e_4 = 1 \iff \begin{cases} \mu(21)\pi + (\mu(121) + \mu(221))\pi^2 \\ \quad + (\mu(1121) + \mu(1221) + \mu(2121) + \mu(2221))\pi^3 \equiv 0 \pmod{\pi^4}, \\ \mu(12)\pi + (\mu(112) + \mu(212))\pi^2 \\ \quad + (\mu(1112) + \mu(1212) + \mu(2112) + \mu(2212))\pi^3 \equiv 0 \pmod{\pi^4}. \end{cases}$$

As in case of e_3 , we have $\mu(121) + \mu(221) \equiv \mu(12) \equiv 0 \pmod{p^2}$. Similarly, $\mu(1121) \equiv \mu(1112) \equiv \binom{\mu(12)}{3} \pmod{\Delta(1121)}$ and $\mu(2221) \equiv \binom{\mu(12)}{3} \pmod{\Delta(2221)}$ by (2.2.6). Since $\Delta(1121) \equiv \Delta(2221) \equiv 0 \pmod{p}$, $\mu(1121) + \mu(2221) \equiv \mu(12)/3 \equiv 0 \pmod{p}$. Finally, by shuffle relation, $\mu(1221) + \mu(2121) + \mu(2211) \equiv \mu(2121) + 2\mu(2211) \equiv 0 \pmod{p}$. Thus the first condition is equivalent to $\mu(21)\pi - \mu(2211)\pi^3 \equiv 0 \pmod{\pi^4}$. Similarly, we see that the second condition is equivalent to $\mu(12)\pi - \mu(1122)\pi^3 \equiv 0 \pmod{\pi^4}$ which is same as the first one. Hence, we obtain

$$e_4 = 1 \iff \text{lk}(K_1, K_2) - \mu(1122)\pi^2 \equiv 0 \pmod{\pi^3}.$$

For case $p = 2$, this is equivalent to the following condition*

$$(4.2.2.4) \quad \begin{cases} \text{lk}(K_1, K_2) \equiv 0 \pmod{8}, \mu(1122) \equiv 0 \pmod{2} \\ \text{or} \\ \text{lk}(K_1, K_2) \equiv 0 \pmod{4}, \text{lk}(K_1, K_2) \not\equiv 0 \pmod{8}, \mu(1122) \equiv 1 \pmod{2}. \end{cases}$$

For example, the Whitehead link $L = K_1 \cup K_2$ satisfies $\text{lk}(K_1, K_2) = 0, \mu(1122) = 1$ and so $e_3 = 1, e_4 = 0$, hence $H_1(M, \mathbf{Z}_p) = \widehat{\mathcal{O}}/\mathfrak{p}^3$. For the 2-bridge link of type (48,37), the latter condition of (4.2.2.4) is satisfied and so $H_1(M, \mathbf{Z}_2) = \mathbf{Z}/2^k\mathbf{Z}, k \geq 4$.

5. Iwasawa type formulas for the p -homology groups of p^m -fold cyclic branched covers

5.1. *Asymptotic formula for the p -homology of p^m -fold covers.* For $m \geq 1$, let M_m be the completion of the p^m -fold subcover of X_∞ over X_L so that M_m is a p^m -fold cyclic cover of S^3 branched along L . In this last Section, we are concerned with the asymptotic behavior of the order of $H_1(M_m, \mathbf{Z}_p)$ as $m \rightarrow \infty$ using the standard argument in Iwasawa theory. As in Section 4, we start again with the following isomorphisms

$$(5.1.1) \quad \widehat{A}_L^{\text{red}} \simeq (H_1(X_\infty, \mathbf{Z}) \otimes_\Lambda \widehat{\Lambda}) \oplus \widehat{\Lambda},$$

$$(5.1.2) \quad H_1(M_m, \mathbf{Z}_p) \simeq (H_1(X_\infty, \mathbf{Z}) \otimes_\Lambda \widehat{\Lambda}) / (\nu_{p^m}(1 + X)).$$

From these, we get immediately an extension of a theorem of M. Dellomo [D] for a knot.

Proposition 5.1.3 *For a link L whose p -adic Milnor invariants are all zero, for example a boundary link, we have $H_1(M_m, \mathbf{Z}_p) = \mathbf{Z}_p^{(p^m-1)(n-1)}$ for $m \geq 1$. In particular, $H_1(M_m, \mathbf{Z}_p) = 0$ for $m \geq 1$ if L is a knot.*

Proof. In fact, $\widehat{A}_L^{\text{red}} = \widehat{\Lambda}^n$ for such a link L by Corollary 3.2.5. Hence $H_1(X_\infty, \mathbf{Z}) \otimes_\Lambda \widehat{\Lambda} = \widehat{\Lambda}^{n-1}$ by (5.1.1) and so $H_1(M_m, \mathbf{Z}_p) = \mathbf{Z}_p^{(p^m-1)(n-1)}$ by (5.1.2). $\square$

In the following, we assume $n \geq 2$. By (5.1.1), the 0-th elementary ideal $E_0(H_1(X_\infty, \mathbf{Z}) \otimes_\Lambda \widehat{\Lambda})$ over $\widehat{\Lambda}$ is same as the 1st completed Alexander ideal $\widehat{E}_1(L)$ (3.1.5). Note that the 1st p -adic Alexander series $\widehat{\Delta}_1(L)$ is given as the greatest common divisors of all $n-1$ minors of the reduced p -adic Traldi matrix and so it is written by the form

$$\widehat{\Delta}_1(L) = X^{n-1} \cdot \widehat{\nabla}_L$$

*K. Murasugi informed us of this condition and examples which are obtained by the relation between the Alexander polynomial and Milnor invariants [Mu].

where we call $\widehat{\nabla}_L$ the p -adic Hosokawa series of L . Then by (5.1.2), we have the following formula on the order $|H_1(M_m, \mathbf{Z}_p)|$ which is seen as the p -primary part of the well known formula by S. Kinoshita and F. Hosokawa [KT] (See also [MM]). Here we interpret $|H_1(M_m, \mathbf{Z}_p)| = 0$ to mean $H_1(M_m, \mathbf{Z}_p)$ is infinite.

Proposition 5.1.4. $|H_1(M_m, \mathbf{Z}_p)| = p^{m(n-1)} \prod_{\substack{\zeta^{p^m}=1 \\ \zeta \neq 1}} |\widehat{\nabla}_L(\zeta - 1)|_p^{-1}.$

Now, we assume $H_1(M_m, \mathbf{Z}_p)$ is finite for any m and see the asymptotic behaviour of the order $|H_1(M_m, \mathbf{Z}_p)|$ as $m \rightarrow \infty$. For this, we recall the following standard facts from Iwasawa theory. We call a polynomial $g(X) \in \mathbf{Z}_p[X]$ *distinguished* if $g(X) = X^d + a_{d-1}X^{d-1} + \cdots + a_0$, $a_i \equiv 0 \pmod{p}$ for $0 \leq i \leq d-1$.

Lemma 5.1.5 (p -adic Weierstrass preparation theorem [W, Theorem 7.3]). *A non-zero element $f(X) \in \widehat{\Lambda}$ is written uniquely as*

$$f(X) = p^\mu g(X) u(X)$$

where μ is a non-negative integer, $g(X)$ is a distinguished polynomial and $u(X) \in \widehat{\Lambda}^\times$.

Lemma 5.1.6 ([W, Theorem 7.14]). *Let $f(X) \in \widehat{\Lambda}$ and assume $f(\zeta - 1) \neq 0$ for any primitive p^m -th root ζ of 1 for $k \geq 1$. Write $f(X) = p^\mu g(X) u(X)$ according to Lemma 5.5 and define λ by the degree of $g(X)$. Then there is an integer ν independent of k such that we have the equality*

$$\text{ord}_p\left(\prod_{\substack{\zeta^{p^m}=1 \\ \zeta \neq 1}} f(\zeta - 1)\right) = \lambda m + \mu p^m + \nu$$

for sufficiently large m .

For the convenience of the reader, we include herewith a proof of Lemma 5.1.6.

Proof of Lemma 5.1.6. Since $u(x) \in \widehat{\Lambda}^\times$, we have

$$\text{ord}_p\left(\prod_{\substack{\zeta^{p^m}=1 \\ \zeta \neq 1}} f(\zeta - 1)\right) = (p^m - 1)\mu + \text{ord}_p\left(\prod_{\substack{\zeta^{p^m}=1 \\ \zeta \neq 1}} g(\zeta - 1)\right).$$

Write $g(X) = X^\lambda + a_{\lambda-1}X^{\lambda-1} + \cdots + a_0$, $a_i \equiv 0 \pmod{p}$ ($0 \leq i \leq \lambda - 1$). For a primitive p^l -th root ζ of 1 ($1 \leq l \leq k$), one has the equality $(p) = (\zeta - 1)^{\phi(p^l)}$ of ideals of $\mathbf{Z}[\zeta]$ and so $\text{ord}_p((\zeta - 1)^\lambda) = \frac{\lambda}{\phi(p^l)}$ where $\phi(x)$ is the Euler function. Therefore, if l is large enough,

$\text{ord}_p((\zeta - 1)^\lambda) < \text{ord}_p(a_i(\zeta - 1)^i)$ for $0 \leq i \leq \lambda - 1$ and so $\text{ord}_p(g(\zeta - 1)) = \text{ord}_p((\zeta - 1)^\lambda)$. Hence, there is a constant C independent of k such that for sufficiently large m , we have

$$\text{ord}_p\left(\prod_{\substack{\zeta^{p^m}=1 \\ \zeta \neq 1}} g(\zeta - 1)\right) = \text{ord}_p\left(\prod_{\substack{\zeta^{p^m}=1 \\ \zeta \neq 1}} (\zeta - 1)^\lambda\right) + C = \text{ord}_p(p^{m\lambda}) + C = \lambda m + C. \quad \square$$

To apply Lemma 5.1.6 to $\widehat{\nabla}_L$, write

$$\widehat{\nabla}_L = p^{\mu(L;p)} g(L;p) u(L;p)$$

where $\mu(L;p)$ is a nonnegative integer, $g(L;p)$ is a distinguished polynomial and $u(L;p) \in \widehat{\Lambda}^\times$ according to Lemma 5.1.5 and set $\lambda(L;p) = \deg(g(L;p))$. Then Proposition 5.1.4 and Lemma 5.1.6 yield the following

Theorem 5.1.7. *Notation and assumption being as above, there is a constant $\nu(L;p)$ depending only on L and p such that we have*

$$\text{ord}_p(|H_1(M_m, \mathbf{Z}_p)|) = (n - 1 + \lambda(L;p))m + \mu(L;p)p^m + \nu(L;p)$$

for sufficiently large m .

We call the invariants $\lambda(L;p), \mu(L;p)$ the *Iwasawa λ, μ -invariants* of L with respect to p respectively after the model of the Iwasawa invariants in the theory of $\mathbf{Z}_p$ -extensions [Iw].

5.2. Examples.

1. Let L be the Whitehead link. We then have

$$\widehat{T}_L = \begin{pmatrix} X_1 X_2^2 & -X_1^2 X_2 \\ X_1^2 X_2 & -X_1 X_2^2 \end{pmatrix}$$

and $\widehat{\nabla}_L = X^2$. Hence, we have $\lambda(L;p) = 2, \mu(L;p) = 0$ and $\text{ord}_p(H_1(M_m, \mathbf{Z}_p)) = 3m$ for $m \geq 1$.

2. Let $L = K_1 \cup K_2 \cup K_3$ be the Borromean rings so that we can take $y_1 = [x_3, x_2], y_2 = [x_3, x_1], y_3 = [x_1, x_2]$. Then we can compute all Milnor number needed to get the reduced Traldi matrix

$$\widehat{T}_L^{\text{red}} = \begin{pmatrix} X + X^2 & -X^2 & -X \\ -X & X + X^2 & -X^2 \\ -X^2 & -X & X + X^2 \end{pmatrix}$$

and so $\widehat{\nabla}_L = 1 + X + X^2$. Hence, we have $\lambda(L;p) = \mu(L;p) = 0$ and $\text{ord}_p(H_1(M_m, \mathbf{Z}_p)) = 2m$ for $m \geq 1$.

REFERENCES

- [A] D. Anick, *Inert sets and the Lie algebra associated to a group*, J. Algebra **111** (1987), 154–165.
- [Da] J. Davis, *The homology of cyclic branched covers of S^3* , Math. Ann. **301** (1995), 507–518.
- [De] M. Dellomo, *On the inverse limit of the branched cyclic covers associated with a knot*, J. Pure Appl. Algebra **40** (1986), 15–26.
- [F] R.H. Fox, *Free differential calculus. I: Derivation in the free group ring*, Ann. of Math **57** (1953), 547–560.
- [H] J.A. Hillman, *Algebraic invariants of links*, Series on Knots and Everything, **32**, World Scientific Publishing Co, 2002.
- [HS] J. Hillman, M. Sakuma, *On the homology of finite abelian coverings of links*, Canad. Math. Bull. **40** (1997), 309–315.
- [HK] F. Hosokawa, S. Kinoshita, *On the homology group of branched cyclic covering spaces of links*, Osaka Math. J. **12** (1960), 331–355.
- [Ih] Y. Ihara, *On Galois representations arising from towers of coverings of $\mathbf{P}^1 \setminus \{0, 1, \infty\}$* , Invent. Math. **86** (1986), 427–459.
- [Iw] K. Iwasawa, *On Γ -extensions of algebraic number fields*, Bull. Amer. Math. Soc. **65** (1959), 183–226.
- [Ko] H. Koch, *Galoissche Theorie der p -Erweiterungen*, Springer, Berlin-New York; VEB Deutscher Verlag der Wissenschaften, Berlin, 1970.
- [L] J.P. Labute, *The Lie algebra associated to the lower central series of a link group and Murasugi’s conjecture*, Proc. Amer. Math. Soc. **109** (1990), 951–956.
- [Mas] W. Massey, *Completion of link modules*, Duke Math. J. **47** (1980), 399–420.
- [Mat] D. Matei, *Fundamental groups of links and arrangements: Characteristic varieties, resonance varieties and finite index subgroups*, Ph.D Thesis, Northeastern University (1999).
- [MM] J.P. Mayberry, K. Murasugi, *Torsion-groups of abelian coverings of links*, Trans. Amer. Math. Soc. **271**, (1982), 143–173.
- [Mi] J. Milnor, *Isotopy of links*, in Algebraic Geometry and Topology, A symposium in honour of S. Lefschetz (edited by R.H. Fox, D.S. Spencer and W. Tucker), Princeton Univ. Press, Princeton (1957), 280–306.
- [Mo1] M. Morishita, *A theory of genera for cyclic coverings of links*, Proc. Japan Academy **77** (2001), 115–118.
- [Mo2] ———, *On certain analogies between knots and primes*, J. Reine Angew. Math. **550** (2002), 141–167.
- [Mo3] ———, *Milnor invariants and Massey products for prime numbers*, Compositio Math. **140** (2004), 69–83.
- [Mo4] ———, *Analogies between knots and primes, 3-manifolds and number fields*, submitted to Proceedings of JAMI conference, Baltimore (2003).
- [Mo5] ———, *Milnor invariants and l -class groups*, submitted (2004).
- [Mu] K. Murasugi, *On Milnor’s invariant for links*, Trans. Amer. Math. Soc. **124** (1966), 94–110.

- [Réd1] L. Rédei, *Arithmetischer Beweis des Satzes über die Anzahl der durch vier teilbaren Invarianten der absoluten Klassengruppe im quadratischen Zahlkörper*, J. Reine Angew. Math. **171** (1934), 55-60.
- [Réd2] ———, *Ein neues zahlentheoretisches Symbol mit Anwendungen auf die Theorie der quadratischen Zahlkörper, I*, J. Reine Angew. Math. **180** (1938), 1-43.
- [Rez1] A. Reznikov, *Three-manifolds class field theory (homology of coverings for a nonvirtually b_1 -positive manifold)*, Selecta Math. New ser. **3** (1997), 361-399.
- [Rez2] ———, *Embedded incompressible surfaces and homology of ramified coverings of three-manifolds*, Selecta Math. New ser. **6** (2000), 1-39.
- [S1] M. Sakuma, *Homology groups of abelian coverings of links*, Math. Seminar Note, Kobe Univ. **7** (1979), 515-530.
- [S2] ———, *Homology of abelian coverings of links and spatial graphs*, Canad. J. Math. **47** (1995), 201-224.
- [Se] J.-P. Serre, *Cohomologie Galoisienne*, Lecture Notes in Mathematics **5**, Springer-Verlag, Berlin - Heidelberg - New York, 1965.
- [Tr] L. Traldi, *Milnor's invariants and the completions of link modules*, Trans. Amer. Math. Soc. **284** (1984), 401-424.
- [Tu] V.G. Turaev, *Milnor's invariants and Massey products*, English transl. J. Soviet Math. **12** (1979), 128-137.
- [W] L. Washington, *Introduction to cyclotomic fields*, Graduate Texts in Mathematics, **83**, Springer-Verlag, New York, 1982.

jonh@maths.usyd.edu.au; matei@ms.u-tokyo.ac.jp; morisita@kenroku.kanazawa-u.ac.jp